

Chalcopyrite-pyrite mineralization associated with orthogneiss, in Lutzow-Holm Bay, Eastern Antarctica

Davaa-Ochir Dashbaatar¹, Yoichi Motoyoshi², Tomokazu Hokada³, Sotaro Baba⁴, Atsushi Kamei⁵, Ippei Kitano⁶,
Prayath Nantasin⁷ and Nugroho Setiawan⁸

¹*Mongolian University of Science and Technology*

^{2,3}*National Institute of Polar Research*

⁴*University of the Ryukyus*

⁵*Shimane University*

⁶*Kyushu University*

⁷*Kasetsart University*

⁸*Gadjah Mada University*

Field and petrographic studies reveal chalcopyrite, or chalcopyrite-pyrite mineralization is found in Lower Proterozoic orthogneiss (tonalite-monzonite-quartz monzonite and metagabbro cumulate) in the Lutzow-Holm Bay area, Eastern Antarctica. Monzonitic rocks are composed of plagioclase-K-feldspar-biotite-stilpnomelane and quartz, and have subduction-related geochemical characteristics, fractionated LREE's and high Sr/Y ratio, whereas metagabbro cumulate comprises clinopyroxene, minor plagioclase, hornblende, quartz, biotite and ilmenite, and is tholeiitic, and magmatically unrelated. The igneous rocks have undergone metamorphism to upper amphibolites facies, shown by garnet intergrowth. Chalcopyrite is intergrown in the metamorphic fabric with all minerals, but some samples reveal chalcopyrite is associated with locally coarser aggregates of biotite. Although speculative, chalcopyrite might have been mobilized and segregated at biotite stability, from the metagabbro cumulate.